

檢測報告

17-0418-007-01-7

第1頁, 共5頁

好健康嚴選

樣品名稱：Super8 好孩子.小兒專用抗氧化益生菌
樣品資訊：有效日期:2020.03.08
樣品狀態：完整包裝
檢測委託單位：麗禧股份有限公司
檢測委託單位地址：台北市大安區樂利路 89 號 3F 前棟

收樣日期：2017/04/18
檢驗日期：2017/04/18 ~ 2017/04/28
報告日期：2017/04/28

檢測項目	結果	定量/偵測極限
西藥成分 195 項	陰性 (未檢出)	如附表

~~~以下空白~~~

### 檢測方法：

1. 西藥成分 195 項：  
實驗室內部方法，以 HPLC 分析

報告簽署人



1. 本測試未涉及檢體抽樣，測試樣品、樣品名稱及樣品資訊係由委託檢驗者提供，實驗室僅負責測試分析，本報告僅對該測試樣品有效。  
The test is exclusive of sampling. Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions. The document refer only to the sample(s) tested.
2. 本報告記載事項僅作為參考資料，不得作為任何商業推銷廣告及訴訟用。  
This document is for reference only and cannot be for any commercial purposes or litigation.
3. 本報告經塗改者無效；本報告內容不得部份複製，僅限整份使用。  
The document will be invalid if altered. The document cannot be reproduced except in full.
4. 檢驗報告僅就委託檢驗者之委託事項提供檢驗結果，至若本產品之合法性，仍應由主管機關依法判斷。  
This document stated test results in accordance with Client's request. The legitimacy of the sample(s) should be judged by the authority.



## 檢測報告

17-0418-007-01-7

第2頁, 共5頁

好健康嚴選

附表

| 序號        | 分析項目                             | LOQ(mg/g) |      | 序號       | 分析項目                   | LOQ(mg/g) |      |
|-----------|----------------------------------|-----------|------|----------|------------------------|-----------|------|
|           |                                  | 液體        | 固體   |          |                        | 液體        | 固體   |
| 類固醇：(10)  |                                  |           |      | 利尿劑：(8)  |                        |           |      |
| 1         | Betamethasone                    | 2         | 10   | 30       | Acetazolamide          | 10        | 50   |
| 2         | Cortisone acetate                | 1         | 5    | 31       | Chlorothiazide         | 0.2       | 1    |
| 3         | Dexamethasone                    | 1         | 5    | 32       | Furosemide             | 0.25      | 1.25 |
| 4         | Hydrocortisone                   | 1         | 5    | 33       | Hydrochlorothiazide    | 0.6       | 3    |
| 5         | Hydrocortisone-21-Acetate        | 1         | 5    | 34       | Indapamide             | 0.75      | 3.75 |
| 6         | Methylprednisolone               | 0.5       | 2.5  | 35       | Metolazone             | 1.75      | 8.75 |
| 7         | Oxymetholone                     | 5         | 25   | 36       | Spironolactone         | 1         | 5    |
| 8         | Prednisolone                     | 0.5       | 2.5  | 37       | Triamterene            | 0.88      | 4.4  |
| 9         | Prednisone                       | 0.2       | 1    | 抗生素：(31) |                        |           |      |
| 10        | Triamcinolone acetonide          | 1         | 5    | 38       | Amoxicillin trihydrate | 1         | 5    |
| 降血脂類：(3)  |                                  |           |      | 39       | Ampicillin trihydrate  | 0.8       | 4    |
| 11        | Benzafibrate                     | 0.06      | 0.3  | 40       | Chloramphenicol        | 0.3       | 1.5  |
| 12        | Clofibrate                       | 0.06      | 0.3  | 41       | Chlortetracycline      | 0.8       | 4    |
| 13        | Gemfibrozil                      | 0.07      | 0.35 | 42       | Erythromycin estolate  | 3.2       | 16   |
| 荷爾蒙類：(14) |                                  |           |      | 43       | Fluconazole            | 0.4       | 2    |
| 14        | 17a-Ethynylestradiol             | 0.164     | 0.82 | 44       | Griseofulvin           | 0.3       | 1.5  |
| 15        | 3,3,5-Triiodothyronine free acid | 0.45      | 2.25 | 45       | Isoniazid              | 0.25      | 1.25 |
| 16        | Carbimazole                      | 0.5       | 2.5  | 46       | Metronidazole          | 1         | 5    |
| 17        | Clomiphene Citrate               | 2.5       | 12.5 | 47       | Nalidixic acid         | 1         | 5    |
| 18        | Diethylstilbestrol               | 1         | 5    | 48       | Nitrofurantoin         | 1         | 5    |
| 19        | Danazol                          | 7.5       | 37.5 | 49       | Nitrofurazone          | 1         | 5    |
| 20        | Estradiol benzoate               | 1         | 5    | 50       | Norfloracin            | 1         | 5    |
| 21        | Estriol                          | 0.25      | 1.25 | 51       | Oxolinic acid          | 1         | 5    |
| 22        | Estrone                          | 0.25      | 1.25 | 52       | Phenazopyridine        | 1         | 5    |
| 23        | Ethynyl-Androstenediol           | 7.5       | 37.5 | 53       | Rifampicin             | 0.8       | 4    |
| 24        | Formestane                       | 15        | 75   | 54       | Sulfadiazine           | 1         | 5    |
| 25        | Methyltestosterone               | 3         | 15   | 55       | Sulfadimethoxine       | 0.3       | 1.5  |
| 26        | Progesterone                     | 2.5       | 12.5 | 56       | Sulfaguanidine         | 0.5       | 2.5  |
| 27        | Testosterone propionate          | 4         | 20   | 57       | Sulfamerazine          | 0.4       | 2    |
| 麻藥拮抗劑：(2) |                                  |           |      | 58       | Sulfamethazine         | 0.45      | 2.25 |
| 28        | Naloxone                         | 0.25      | 1.25 | 59       | Sulfamethizole         | 0.3       | 1.5  |
| 29        | Naltrexone                       | 0.25      | 1.25 | 60       | Sulfamethoxazole       | 0.2       | 1    |

- 本測試未涉及檢體抽樣，測試樣品、樣品名稱及樣品資訊係由委託檢驗者提供，實驗室僅負責測試分析，本報告僅對該測試樣品有效。  
The test is exclusive of sampling. Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions. The document refer only to the sample(s) tested.
- 本報告記載事項僅作為參考資料，不得作為任何商業推銷廣告及訴訟用。  
This document is for reference only and cannot be for any commercial purposes or litigation.
- 本報告經塗改者無效；本報告內容不得部份複製，僅限整份使用。  
The document will be invalid if altered. The document cannot be reproduced except in full.
- 檢驗報告僅就委託檢驗者之委託事項提供檢驗結果，至若本產品之合法性，仍應由主管機關依法判斷。  
This document stated test results in accordance with Client's request. The legitimacy of the sample(s) should be judged by the authority.



## 檢測報告

17-0418-007-01-7

第3頁, 共5頁

好健康嚴選

| 序號               | 分析項目                                     | LOQ(mg/g) |       | 序號                | 分析項目                   | LOQ(mg/g) |       |
|------------------|------------------------------------------|-----------|-------|-------------------|------------------------|-----------|-------|
|                  |                                          | 液體        | 固體    |                   |                        | 液體        | 固體    |
| 61               | Sulfamethoxypyridazine                   | 0.4       | 2     | 92                | Metoclopramide         | 0.6       | 3     |
| 62               | Sulfanilamide                            | 0.4       | 2     | 93                | Oxyphencyclimine       | 1         | 5     |
| 63               | Sulfathiazole                            | 0.5       | 2.5   | 94                | Proprantheline bromide | 0.4       | 2     |
| 64               | Sulfisomidine                            | 1         | 5     | 95                | Ranitidine HCL         | 3.33      | 16.65 |
| 65               | Sulfisoxazole                            | 1         | 5     | <b>精神科類：(14)</b>  |                        |           |       |
| 66               | Tetracycline HCl                         | 0.8       | 4     | 96                | Chlormezanone          | 0.125     | 0.625 |
| 67               | Trimethoprim                             | 0.2       | 1     | 97                | Chlorpromazine HCl     | 0.25      | 1.25  |
| 68               | Econazole Nitrate                        | 0.588     | 2.94  | 98                | Clordiazepoxide HCl    | 0.25      | 1.25  |
| <b>糖尿病用藥：(7)</b> |                                          |           |       | 99                | Diazepam               | 0.25      | 1.25  |
| 69               | Acetohexamide                            | 0.5       | 2.5   | 100               | Diphenylhydantoin      | 0.125     | 0.625 |
| 70               | Chlorpropamide                           | 0.25      | 1.25  | 101               | Flunitrazepam(FM2)     | 0.155     | 0.775 |
| 71               | Glybenclamide                            | 0.25      | 1.25  | 102               | Fluoxetine             | 0.5       | 2.5   |
| 72               | Metformin                                | 0.5       | 2.5   | 103               | Melatonin              | 0.4       | 2     |
| 73               | Phenformin hydrochloride                 | 0.5       | 2.5   | 104               | Meprobamate            | 10        | 50    |
| 74               | Tolazamide                               | 0.125     | 0.625 | 105               | Oxazolam               | 2         | 10    |
| 75               | Tolbutamide                              | 0.25      | 1.25  | 106               | Phenobarbital          | 0.25      | 1.25  |
| <b>抗組織胺類：(8)</b> |                                          |           |       | 107               | Primidone              | 0.25      | 1.25  |
| 76               | Brompheniramine maleate                  | 0.125     | 0.625 | 108               | Sulpiride              | 1         | 5     |
| 77               | Carbinoxamine maleate                    | 0.125     | 0.625 | 109               | Thioridazine           | 0.5       | 2.5   |
| 78               | Chlorpheniramine maleate                 | 0.5       | 2.5   | <b>解熱止痛劑：(27)</b> |                        |           |       |
| 79               | Cinnarizine                              | 0.125     | 0.625 | 110               | Acemetacin             | 0.2       | 1     |
| 80               | Diphenhydramine                          | 0.53      | 2.65  | 111               | Acetaminophen          | 0.25      | 1.25  |
| 81               | Diphenidol                               | 0.14      | 0.7   | 112               | Acetylsalicylic acid   | 1         | 5     |
| 82               | Ketotifen Fumarate                       | 0.25      | 1.25  | 113               | Aminopyrine            | 0.3       | 1.5   |
| 83               | Triprolidine                             | 0.5       | 2.5   | 114               | Benzydamine            | 0.2       | 1     |
| <b>腸胃類：(11)</b>  |                                          |           |       | 115               | Bucetin                | 0.25      | 1.25  |
| 84               | Atropine                                 | 0.4       | 2     | 116               | Bufexamac              | 0.1       | 0.5   |
| 85               | Bisacodyl                                | 0.4       | 2     | 117               | Diclofenac sodium      | 0.1       | 0.5   |
| 86               | Buscopan<br>(scopolamine n-butylbromide) | 1         | 5     | 118               | Dipyrone               | 2         | 10    |
| 87               | Camylofine                               | 1         | 5     | 119               | Ethoxybenzamide        | 0.1       | 0.5   |
| 88               | Cimetidine                               | 0.4       | 2     | 120               | Fenbufen               | 0.25      | 1.25  |
| 89               | DL-Homatropine                           | 0.4       | 2     | 121               | Flufenamic acid        | 0.2       | 1     |
| 90               | Ethaverine                               | 0.3       | 1.5   | 122               | Flurbiprofen           | 0.2       | 1     |
| 91               | Flopropione                              | 0.3       | 1.5   | 123               | Ibuprofen              | 0.2       | 1     |

1. 本測試未涉及檢體抽樣，測試樣品、樣品名稱及樣品資訊係由委託檢驗者提供，實驗室僅負責測試分析，本報告僅對該測試樣品有效。  
The test is exclusive of sampling. Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions. The document refer only to the sample(s) tested.
2. 本報告記載事項僅作為參考資料，不得作為任何商業推銷廣告及訴訟用。  
This document is for reference only and cannot be for any commercial purposes or litigation.
3. 本報告經塗改者無效；本報告內容不得部份複製，僅限整份使用。  
The document will be invalid if altered. The document cannot be reproduced except in full.
4. 檢驗報告僅就委託檢驗者之委託事項提供檢驗結果，至若本產品之合法性，仍應由主管機關依法判斷。  
This document stated test results in accordance with Client's request. The legitimacy of the sample(s) should be judged by the authority.



## 檢測報告

17-0418-007-01-7

第4頁, 共5頁

好健康嚴選

| 序號         | 分析項目             | LOQ(mg/g) |       | 序號              | 分析項目                                 | LOQ(mg/g) |       |
|------------|------------------|-----------|-------|-----------------|--------------------------------------|-----------|-------|
|            |                  | 液體        | 固體    |                 |                                      | 液體        | 固體    |
| 124        | Indomethacin     | 0.15      | 0.75  | 155             | Tetracaine                           | 0.5       | 2.5   |
| 125        | Ketoprofen       | 0.15      | 0.75  | 痛風治療類：(5)       |                                      |           |       |
| 126        | Mefenamic acid   | 0.25      | 1.25  | 156             | Allopurinol                          | 1         | 5     |
| 127        | Methylsalicylate | 0.04      | 0.2   | 157             | Benzbromarone                        | 0.25      | 1.25  |
| 128        | Naproxen         | 0.4       | 2     | 158             | Colchicine                           | 0.25      | 1.25  |
| 129        | Niflumic acid    | 0.35      | 1.75  | 159             | Probenecid                           | 0.25      | 1.25  |
| 130        | Oxyphenbutazone  | 0.6       | 3     | 160             | Sulfinpyrazone                       | 0.25      | 1.25  |
| 131        | Phenacetin       | 0.2       | 1     |                 |                                      |           |       |
| 132        | Phenylbutazone   | 0.5       | 2.5   | 成癮性麻醉劑、禁用藥物：(7) |                                      |           |       |
| 133        | Piroxicam        | 0.4       | 2     | 161             | Amphetamine(安非他命)                    | 1         | 5     |
| 134        | Salicylamide     | 0.1       | 0.5   | 162             | Fenfluramine                         | 1         | 5     |
| 135        | Salicylic acid   | 0.385     | 1.925 | 163             | Heroin(海洛因)                          | 1         | 5     |
| 136        | Sulindac         | 0.1       | 0.5   | 164             | Ketamine(K他命)                        | 1         | 5     |
| 心臟血管類：(13) |                  |           |       | 165             | Methamphetamine(甲基安非他命)              | 1         | 5     |
| 137        | Atenolol         | 1         | 5     | 166             | Phentermine                          | 0.25      | 1.25  |
| 138        | Diltiazem        | 0.25      | 1.25  | 167             | Secobarbital                         | 0.66      | 3.3   |
| 139        | Dipyridamole     | 1         | 5     | 其他：(6)          |                                      |           |       |
| 140        | Fenofibrate      | 0.37      | 1.85  | 168             | Chlorhexidine                        | 0.4       | 2     |
| 141        | Hydralazine HCl  | 2         | 10    | 169             | Methimazole(抗甲狀腺藥物)                  | 1.3       | 6.5   |
| 142        | Methyldopa       | 0.25      | 1.25  | 170             | Mixed alkyltrimethylammonium bromide | 0.8       | 4     |
| 143        | Minoxidil        | 0.25      | 1.25  | 171             | Nicotine(尼古丁)                        | 0.4       | 2     |
| 144        | Nifedipine       | 0.5       | 2.5   | 172             | Podophyllotoxin(鬼臼毒素)                | 0.2       | 1     |
| 145        | Pentoxifylline   | 0.25      | 1.25  | 173             | Santonin(山道年)-驅蟲劑                    | 2.3       | 11.5  |
| 146        | Prazosin         | 1         | 5     | 神經興奮劑：(6)       |                                      |           |       |
| 147        | Propranolol      | 1         | 5     | 174             | Caffeine(咖啡因)                        | 0.25      | 1.25  |
| 148        | Reserpine(蛇根鹼)   | 1         | 5     | 175             | Ephedrine(麻黃素)                       | 2.5       | 12.5  |
| 149        | Phentolamine     | 0.25      | 1.25  | 176             | Methylephedrine                      | 1         | 5     |
| 局部麻醉類：(6)  |                  |           |       | 177             | Phenylephrine                        | 2         | 10    |
| 150        | Benzocaine       | 0.125     | 0.625 | 178             | Phenylpropanolamine                  | 4         | 20    |
| 151        | Dibucaine        | 0.593     | 2.965 | 179             | Theobromine(可可鹼)                     | 1         | 5     |
| 152        | Lidocaine        | 0.25      | 1.25  | 肌肉鬆弛劑：(5)       |                                      |           |       |
| 153        | Oxethazaine      | 0.25      | 1.25  | 180             | Carisoprodol                         | 10        | 50    |
| 154        | Procaine         | 5.81      | 29.05 | 181             | Chlorzoxazone                        | 0.625     | 3.125 |

- 本測試未涉及檢體抽樣，測試樣品、樣品名稱及樣品資訊係由委託檢驗者提供，實驗室僅負責測試分析，本報告僅對該測試樣品有效。  
The test is exclusive of sampling. Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions. The document refer only to the sample(s) tested.
- 本報告記載事項僅作為參考資料，不得作為任何商業推銷廣告及訴訟用。  
This document is for reference only and cannot be for any commercial purposes or litigation.
- 本報告塗改者無效；本報告內容不得部份複製，僅限整份使用。  
The document will be invalid if altered. The document cannot be reproduced except in full.
- 檢驗報告僅就委託檢驗者之委託事項提供檢驗結果，至若本產品之合法性，仍應由主管機關依法判斷。  
This document stated test results in accordance with Client's request. The legitimacy of the sample(s) should be judged by the authority.



## 檢測報告

17-0418-007-01-7

第5頁, 共5頁

好健康嚴選

| 序號         | 分析項目                                                      | LOQ(mg/g) |        | 序號         | 分析項目             | LOQ(mg/g) |       |
|------------|-----------------------------------------------------------|-----------|--------|------------|------------------|-----------|-------|
|            |                                                           | 液體        | 固體     |            |                  | 液體        | 固體    |
| 182        | Flavoxate                                                 | 0.25      | 1.25   | 鎮咳、祛痰劑：(7) |                  |           |       |
| 183        | Mephenesin                                                | 0.125     | 0.625  | 189        | Becantex         | 8         | 40    |
| 184        | Orphenadrine citrate                                      | 0.25      | 1.25   | 190        | Bromhexine       | 2         | 10    |
| 支氣管擴張劑：(4) |                                                           |           |        | 191        | Cloperastine     | 0.5       | 2.5   |
| 185        | Aminophylline                                             | 3.375     | 16.875 | 192        | Dextromethorphan | 1         | 5     |
|            | (Theophylline)                                            |           |        | 193        | Guaifenesin      | 0.125     | 0.625 |
| 186        | Dihydroxypropyltheophylline<br>(Diprophylline、Dyphylline) | 0.25      | 1.25   | 194        | Noscapine HCl    | 1         | 5     |
| 187        | Fenoterol                                                 | 4.25      | 21.25  | 195        | Papaverine(罌粟鹼)  | 0.5       | 2.5   |
| 188        | Salbutamol                                                | 1         | 5      |            |                  |           |       |

- 本測試未涉及檢體抽樣，測試樣品、樣品名稱及樣品資訊係由委託檢驗者提供，實驗室僅負責測試分析，本報告僅對該測試樣品有效。  
The test is exclusive of sampling. Any holder of this document is advised that information contained hereon reflects the Company's findings at the time of its intervention only and within the limits of Client's instructions. The document refer only to the sample(s) tested.
- 本報告記載事項僅作為參考資料，不得作為任何商業推銷廣告及訴訟用。  
This document is for reference only and cannot be for any commercial purposes or litigation.
- 本報告經塗改者無效；本報告內容不得部份複製，僅限整份使用。  
The document will be invalid if altered. The document cannot be reproduced except in full.
- 檢驗報告僅就委託檢驗者之委託事項提供檢驗結果，至若本產品之合法性，仍應由主管機關依法判斷。  
This document stated test results in accordance with Client's request. The legitimacy of the sample(s) should be judged by the authority.